

AC Joint Reconstruction Physical Therapy Protocol

Phase 1: Protection and Range of Motion (Weeks 0–4)

Sling Usage:

- Sling with abduction pillow (discontinue pillow at Week 4).

Restrictions:

- No active range of motion (AROM) until Week 8.

Exercises:

- Pendulums
- Scapular Retractions: Avoid shoulder shrugs.
- Active ROM: Elbow, wrist, and hand.
- Passive ROM:
 - Shoulder flexion and abduction to 90°.
 - External rotation (ER) and internal rotation (IR) unrestricted.

Phase 2: Early Strengthening (Weeks 4–6)

Key Changes:

- Discontinue abduction pillow.

Exercises:

- Isometrics: Begin for ER and IR.
 - Assisted Active Range of Motion (AAROM):
 - Flexion and abduction to 90°.
 - Full ER and IR.
-

Phase 3: Strength and Mobility (Weeks 6–8)

Key Changes:

- Discontinue sling.
- Progress to full AAROM as tolerated.

Exercises:

- Unrestricted Isometrics: Rotator cuff (RTC) and periscapular muscles.

“Dr. Trevor” Stefanski, M.D.
Minimally Invasive Joint Specialist
DrTrevor.com/PT for a printable copy

Innovating **BIG RESULTS**
Through Small Incisions

- Concentric and Eccentric Exercises: Begin at Week 8.
-

Advanced Sports-Specific Training (Week 16 Onwards)

Key Focus:

- Advanced weight-bearing exercises.
- Plyometric exercises.
- Sports-specific drills.

Phone: 513-232-2663
Fax: 513-985-2580